DRAFT VERSION APRIL 24, 2013
Preprint typeset using LATEX style emulateapj v. 11/10/09

arXiv:1304.6137v1 [astro-ph.HE] 23 Apr 2013

THE FERMI BUBBLES AS STARBURST WIND TERMINATION SHOCKS

BRIAN C. LACKI[1,2]

Draft version April 24, 2013

ABSTRACT

The enhanced star formation in the inner 100 pc of the Galaxy launches a superwind, reaching ~ 1600 km s^{-1} for M82-like parameters. The ram pressure of the wind is very low compared to more powerful starburst winds. I show that halo gas stops the wind a few kpc from the Galactic Center. I argue that the termination shock accelerates cosmic rays, and that the resulting Inverse Compton γ-rays are visible as the Fermi Bubbles. The Bubbles are thus wind bubbles, which the starburst can inflate within 10 Myr. They remain in steady state as long as the starburst lasts. The shock may accelerate PeV electrons and ultra-high energy protons. The Bubbles may be analogs of galactic wind termination shocks in the intergalactic medium. I discuss the advantages and problems of this model.

Subject headings: Galaxy: center — galaxies: starburst — ISM: jets and outflows — gamma rays: galaxies — cosmic rays

1. INTRODUCTION

Powerful outflows are ever present in starburst regions ($\Sigma_{\rm SFR} \gtrsim 0.1$ M$_\odot$ yr^{-1} kpc^{-2}), blasting out at speeds $\gg$ 100 km s^{-1} (Chevalier & Clegg 1985 (CC85); Heckman et al. 1990; Strickland & Heckman 2009). Winds erupt in these intense regions when superbubbles and supernova remnants overlap as the star-formation density increases (McKee & Ostriker 1977). The hot plasma fills starbursts and blows out in a sound-crossing time, continuously replenished by massive stars. Even in normal galaxies like the Milky Way, cosmic rays (CRs) can drive winds once they diffuse far enough from the galactic plane (Breitschwerdt et al. 1991). But what happens to the wind after it escapes these central regions?

In the CC85 model, the wind rapidly accelerates as it passes the sonic point, reaching an asymptotic speed $v_\infty \approx$ 1600 km s^{-1} if supernova energy thermalizes efficiently and the mass loading is small as in M82 (Strickland & Heckman 2009). As the wind expands adiabatically out beyond the confines of the starburst, particle random motions in the wind slow down. This applies both to the wind plasma itself and CRs within the winds (Völk et al. 1996). The random kinetic energy of the wind is converted into bulk kinetic energy, pushing the wind plasma like a piston.

The wind usually escapes the galaxy, growing ever more dilute. Eventually, it becomes so rarefied that the ram pressure equals whatever external pressure there is. At this location, a termination shock stops the wind. These shocks are theorized to host energetic phenomena, such as cosmic ray acceleration. However, the termination shock of starbursts lie in the intergalactic medium.

A prototype for starburst regions and outflows exists in our Galaxy, in the form of the Galactic Center Central Molecular Zone (CMZ). Lying within 100 pc of Sgr A*, the region is filled with gas (Molinari et al. 2011), a large star-formation rate density ($\Sigma_{\rm SFR} \approx 2$ M$_\odot$ yr^{-1} kpc^{-2}; Yusef-Zadeh et al. 2009), strong magnetic fields (Crocker et al. 2010), and TeV emission (Aharonian et al. 2006). Evidence for an outflow includes the abnormally weak nonthermal radio and γ-ray emission (Crocker et al. 2011a), suggesting that CRs are advected away from the CMZ (Crocker et al. 2011b).

The Fermi Bubbles are more evidence for powerful phenomena in the Galactic Center (Finkbeiner 2004; Dobler et al. 2010; Su et al. 2010; Ade et al. 2012). These bilobal structures, visible in γ-rays and radio, extend nearly 10 kpc from the Galactic Plane and emanate from the Center. Soft X-ray emission is suggestive of a shock at the Bubble edges (Sofue 2000; Bland-Hawthorn & Cohen 2003; Su et al. 2010). Most theoretical work interprets these bubbles as the result of an outburst from Sgr A* (Cheng et al. 2011; Zubovas et al. 2011; Guo & Mathews 2012). In these models, CR electrons (e^-) shine in radio by synchrotron emission and in γ-rays by Inverse Compton (IC) emission. But, since these losses are rapid, either the e^- are transported rapidly or they must be accelerated in place.

Crocker & Aharonian (2011) proposed an alternative explanation, in which the CMZ powers the Bubbles. In this model, the CMZ has been accelerating CRs for the entire Galaxy's history. The wind advects CR protons a few kpc, where they accumulate for ten billion years. The Bubbles' γ-ray emission is pionic; secondary $e^\pm$ made by pionic interactions emit the radio waves. Since the gas density is low in the Bubbles ($\lesssim 0.01$ cm^{-3}), the CRs must accumulate for ten billion years in order to make enough γ-rays. This model evades the severe adiabatic losses in a CC85-style wind by postulating a slow wind that becomes compressed in the Bubbles. Leptonic starburst models are possible too: Biermann et al. (2010) argued that high energy $e^\pm$ rapidly diffuse from the Center's starburst and are advected into the Bubbles, where they emit IC.

In this Letter, I propose that the Fermi Bubbles are the termination shocks of the Center's starburst wind (Figure 1). Because the CMZ wind is less dense than prototypical starbursts' winds, its ram pressure is quite low. Thus, the wind is stopped by the Galactic halo. These shocks accelerate CRs. I suggest that primary e^- are the sources of the nonthermal emission. Thus, the Fermi Bubbles are a nearby analog of the intergalactic termination shocks of more powerful starbursts (c.f., Jokipii & Morfill 1987; Völk & Zirakashvili 2004).

---

[1] Jansky Fellow of the National Radio Astronomy Observatory
[2] Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA, brianlacki@ias.edu



FIG. 1.— Sketch of the CMZ and its termination shock with respect to the Galaxy. CR e^- and p^+ are accelerated at the shock; the emission from the e^- appears to us as the radio and γ-ray bubbles.

2. PROPERTIES OF THE CMZ WIND AND ITS TERMINATION SHOCK

2.1. *Basic wind properties*

My discussion is informed by the CC85 theory of starburst winds, as amended by Strickland & Heckman (2009) to apply to disk geometries. A hot plasma fills the wind volume, with properties determined by energy and mass injection. The CMZ is a disk with radius $R_{\rm CMZ}$ = 112 pc and height $h_{\rm CMZ}$ = 42 pc, forming stars at a rate SFR $\approx$ 0.1 $M_{\odot}$ yr^{-1} (Crocker et al. 2011b). Mass is injected into the wind at a rate $\dot{M} = 0.117\beta \times {\rm SFR}$ from stellar winds and explosions, and these phenomena heat the wind at a rate $\dot{E} = 2.54 \times 10^{41} \epsilon_{\rm therm} [{\rm SFR}/(M_{\odot}\ {\rm yr}^{-1})]$erg sec^{-1}.[3] I take $\beta = 2$ and $\epsilon_{\rm therm} = 0.75$, as for M82 (Strickland & Heckman 2009). Then the central density of the wind is given by

$$n_c = \frac{0.592\sqrt{\dot{M}^3/\dot{E}}}{R_{\rm CMZ}^2 (1 + \frac{2h_{\rm CMZ}}{R_{\rm CMZ}})\mu m_H} = 0.037\ {\rm cm}^{-3} \left(\frac{\rm SFR}{0.1\ M_{\odot}\ {\rm yr}^{-1}}\right) \tag{1}$$

and the central temperature is

$$T_c = \frac{(\gamma - 1)\mu m_H \dot{E}}{\gamma k_B \dot{M}} = 3.7 \times 10^7\ {\rm K}, \tag{2}$$

for a mean molecular weight $\mu = 0.59$ and adiabatic index $\gamma = 5/3$.

In the CC85 model, the wind flows out of the starburst at a speed $v_{\infty} = \sqrt{2\dot{E}/\dot{M}} = 1600$ km s^{-1}. The escape speed from the CMZ is only 800 – 1000 km s^{-1} (Breitschwerdt et al. 1991; Launhardt et al. 2002), so the wind escapes.[4] Although CC85 assume spherical symmetry, actual starburst winds are observed to flow in cones. I therefore assume the outflow cone fills a solid angle Ω. The density of the outflow is $\rho = \dot{M}/(\Omega v_{\infty} r^2)$, and the ram pressure is $P_{\rm ram} = \rho v_{\infty}^2/2$.

2.2. *The Termination Shock*

The termination shock occurs when the ram pressure is equal to the external pressure. Supposing the external pressure is entirely thermal, $P_{\rm ext} = n_{\rm ext} k_B T_{\rm ext}$, the termination shock distance is

$$R_t = \sqrt{\frac{\dot{M} v_{\infty}}{2\Omega n_{\rm ext} k_B T_{\rm ext}}}. \tag{3}$$

The state of the gas several kpc into the Galactic halo, and thus the location of the CMZ termination shock, is poorly known. I take $P_{\rm ext}/k_B = 1000$ K cm^{-3}, consistent with loose constraints from Galactic absorption line features (e.g., Savage et al. 2003; Yao & Wang 2005; Hsu et al. 2011) and hydrodynamics (e.g., Spitzer 1956; Wolfire et al. 1995; Fang et al. 2013). R_t depends weakly on the external pressure.

For the CMZ wind,

$$R_t = 5.4\ {\rm kpc} \left(\frac{\rm SFR}{0.1\ M_{\odot}\ {\rm yr}^{-1}}\right)^{1/2} \left(\frac{\Omega}{\pi}\right)^{-1/2} \left(\frac{P_{\rm ext}/k_B}{10^3\ {\rm K\ cm}^{-3}}\right)^{-1/2}, \tag{4}$$

roughly the size of the Fermi Bubbles. For SFRs of 3 to 200 $M_{\odot}$ yr^{-1} (e.g., NGC 253, M82, or Arp 220), the termination shock is instead firmly in the realm of circumgalactic or even intergalactic distances.

The termination shock is a steady-state feature, present even if the starburst wind has always been active. The wind just inside the shock is in pressure equilibrium with the gas outside. In contrast, many models of the Fermi Bubbles propose that the shock is transient, resulting from a recent burst of energy injection, and the shock is driven outwards by an overpressure.

By the time the wind reaches these distances, it is very cold. Outside of the starburst, where there is no energy injection into the wind, the temperature of the wind scales as $T \propto \rho^{\gamma-1} \propto \rho^{2/3}$; likewise the energy density of relativistic CRs scales as $\rho^{1/3}$. After scaling to the central density and temperature,

$$T(R) \approx 1.7 \times 10^5\ {\rm K} \left(\frac{R}{5\ {\rm kpc}}\right)^{-4/3} \left(\frac{\Omega}{\pi}\right)^{-2/3} \tag{5}$$

$$\frac{U_{\rm CR}(R)}{U_{\rm CR}(0)} \approx 0.069 \left(\frac{R}{6\ {\rm kpc}}\right)^{-2/3} \left(\frac{\Omega}{\pi}\right)^{-1/3}. \tag{6}$$

The adiabatic cooling time $(3/4)R/v_{\infty} \approx 2.8$ Myr $(R/6\ {\rm kpc})$ is comparable to the radiative cooling time for high ($\sim 3Z_{\odot}$) metallicity gas at these distances (Sutherland & Dopita 1993).

2.3. *Evolution of the Fermi Bubbles*

Some authors previously proposed, not a permanent CMZ wind, but a wind from a brief CMZ starburst 15 Myr ago; the shock front from the wind could appear as radio/X-ray bubbles (Sofue 2000; Bland-Hawthorn & Cohen 2003). If the starburst wind just turned on, how long does it take to reach steady state and inflate the Bubbles? Suppose that the medium surrounding the CMZ is uniform, with density $\rho_{\rm ext}$ and pressure $P_{\rm ext} = \rho_{\rm ext}\sigma^2/2$. The wind turns on, and starts sweeping up this medium at a shock. Once it reaches a radius $R_{\rm Sed} \approx \sqrt{3\dot{M}/(\Omega v_\infty \rho_{\rm ext})}$, the mass of swept up material is greater than the mass in the wind, and the shock slows down – much like a supernova remnant does, except there is continuous mass injection. As long as $\sqrt{3}\sigma < v_\infty$, $R_{\rm Sed} < R_t$; for the conditions I have been assuming, $R_{\rm Sed} \approx 1$ kpc.

---

[3] I have converted the values in Strickland & Heckman (2009) to a Salpeter IMF running from 0.1 $M_{\odot}$ to 100 $M_{\odot}$.

[4] Carretti et al. (2013) argue that the radio "spurs" trace CMZ features if the outflow has a speed of 1000 km s^{-1}.

After this point, I estimate the shock speed by assuming that the Bubbles are adiabatic, and that the kinetic energy of the swept up material is a fraction κ of the injected mechanical power. The radius of the shock is $dR_s/dt = \sqrt{6\kappa\dot{E}t/(\rho_{\rm ext}\Omega R_s^3)}$, giving $t = R^{5/3}[(3/50)\rho_{\rm ext}/(\dot{E}\kappa)]^{1/3}$. It takes

$$t_{\rm inflate} = \left[\frac{9}{80000}\frac{\dot{M}^5 v_\infty^5 \rho_{\rm ext}^2}{\kappa^2 \dot{E}^2 \Omega^3 P_{\rm ext}^5}\right]^{1/6} \approx 9\kappa^{-1/3}\ {\rm Myr} \qquad (7)$$

for the shock to reach the location of the termination shock (with $\rho_{\rm ext} = 0.001 m_H$ cm^{-3} and $P_{\rm ext} = 1000$ K cm^{-3}), where the shock expands at a speed $v_s = 0.761\kappa^{1/3}(v_\infty/\sigma)^{1/3}\sigma \approx 230\kappa^{1/3}$ km s^{-1}. The shock initially overshoots R_t in this picture, but then it oscillates back and forth before settling down at R_t.

In my model, $t_{\rm inflate}$ is longer than the minimum time $H/\dot{E} \approx 2$ Myr for the starburst to do enough work to inflate the Bubbles, where $H = \gamma P_{\rm ext}[\Omega R_t^3/3]$ is the enthalpy (c.f., Crocker 2012). The reason is that the swept up material is not simply pushed out of the Bubbles, but accelerated to v_s, requiring more energy. On the other hand, my estimate of $H/\dot{E}$ is ~ 3000 times lower than the 10 Gyr estimated by Crocker (2012). The major discrepancy is caused by three factors. First, the estimated volume of the Fermi Bubbles in this discussion is $\Omega R_t^3/3 \approx 5 \times 10^{66}$ cm^3, whereas Crocker (2012) uses 2×10^{67} cm^3; the latter value is more accurate. Second and more importantly, Crocker (2012) uses an external pressure of $P/k_B \approx 4 \times 10^4$ K cm^{-3}. This pressure is characteristic of the inner Milky Way disk, but the pressure in the halo (and most of the volume) could be much lower. Finally, Crocker (2012) uses the minimum outflow power of $\dot{E} \approx 10^{39}$ erg s^{-1}, from the expected CR injection rate. In any case, it is clear that the CMZ can inflate the Fermi Bubbles.

What if the wind suddenly shuts off? Then the Bubbles collapse at a speed $\sim \sigma$, specifically the sound speed. The collapse time is very long, though, almost 40 Myr. The large stellar (Launhardt et al. 2002) and gas (Molinari et al. 2011) masses in the nuclear bulge are consistent with star-formation over several Gyr. Even if the output power of the CMZ fluctuates on timescales of $\lesssim 10$ Myr, as extragalactic true nuclear starbursts do (Mayya et al. 2004), the termination shock remains roughly in the same place. Thus, the Fermi Bubbles may be a relatively permanent feature of our Galaxy, as Crocker & Aharonian (2011) originally argued.

3. NONTHERMAL EMISSION FROM THE SHOCK

Because of the adiabatic losses, only a small fraction of CR energy density remains as the wind reaches its termination shock, regardless of any additional radiative losses. Yet neither the CR nor the thermal energy disappears during adiabatic expansion; it simply converts into bulk kinetic energy. At the termination shock, that energy is converted back into random particle energy. Most of the energy goes into heating the gas, but an appreciable factor ($\gtrsim 10\%$) is in the form of relativistic particles accelerated at the shock. Thus, termination shocks at the Bubble edges inject relativistic particles *in situ*, circumventing any losses.

3.1. *Estimate of γ-ray emission*

The mechanical power of the wind is just $\dot{E}$. Suppose the shocks convert $\eta = 30\%$ of this power into CR protons, which are accelerated with a E^{-2} spectrum. I define $\Psi = \ln(E_{\rm max}/E_{\rm min})$, where $E_{\rm min}$ and $E_{\rm max}$ are the minimum and maximum energies of CR protons in the spectrum. Then the injection spectrum of CR protons is:

$$E^2 \frac{dQ_p}{dE} \approx 2.9 \times 10^{38}\ {\rm erg\ s^{-1}\ SFR_{0.1}} \eta_{0.3} \Psi_{20}^{-1}, \qquad (8)$$

where ${\rm SFR}_{0.1} = {\rm SFR}/(0.1\ M_\odot\ {\rm yr}^{-1})$, $\eta_{0.3} = \eta/0.3$, and $\Psi_{20} = \Psi/20$. The protons presumably diffuse out, or join a larger-scale Galactic wind, without radiating.

Collisionless shocks also accelerate primary CR electrons, but $\gtrsim$ GeV electrons are thought to carry little of the energy. In the test particle approximation to diffusive shock acceleration, at energies $\gtrsim m_p c^2$, the ratio of injected electrons to injected protons is $\tilde{\delta} = (m_p/m_e)^{(p-1)/2}$ (Bell 1978). Thus, for $p = 2.0$, we have $\tilde{\delta} \approx 43$:

$$E^2 \frac{dQ_e}{dE} \approx 6.7 \times 10^{36}\ {\rm erg\ s^{-1}\ SFR_{0.1}} \eta_{0.3} \Psi_{20}^{-1}. \qquad (9)$$

Radiative losses for electrons are fast near the Galaxy, and the IC cooling time for an electron is

$$t_{\rm IC} = 3.1\ {\rm Myr} \left(\frac{E}{100\ {\rm GeV}}\right)^{-1} \left(\frac{U_{\rm rad}}{1\ {\rm eV\ cm^{-3}}}\right)^{-1}. \qquad (10)$$

I assume that a fraction $f_{\rm IC} \approx 1$ of the power goes into IC, with the rest mostly going into synchrotron. In the Thomson limit, the energy of an IC upscattered photon goes as $E_\gamma \propto E_e^2$, where E_e is the relativistic electron energy. Thus, each dex of electron energy is stretched out into 2 dex of γ-ray energy, and we have $E^2 dL/dE \approx (f_{\rm IC}/2) E^2 dQ_e/dE$. The average distance of the Bubbles is 10 kpc, and they cover 0.8 sr (Su et al. 2010). So the flux from the Fermi Bubbles at Earth is

$$E^2 \frac{dN_{\rm IC}}{dE} \approx 220\ {\rm eV\ cm^{-2}\ sec^{-1}\ sr^{-1}}\ f_{\rm IC} {\rm SFR}_{0.1} \eta_{0.3} \Psi_{20}^{-1}. \qquad (11)$$

This matches the γ-ray flux from the Fermi Bubbles (Su et al. 2010).

3.2. *Magnetic fields*

Synchrotron microwaves from the Bubbles demonstrate that magnetic fields are present. Its 23 GHz brightness temperature is ~ 50 μKelv at $|b| \lesssim 30°$ (Dobler 2012a; Ade et al. 2012). From minimum energy arguments (Beck & Krause 2005) and γ-ray to radio ratios (Dobler 2012b; Hooper & Slatyer 2013), the magnetic fields are roughly 4 – 6 μG. The radio "spurs" may have higher magnetic fields, ~ 15 μG (Jones et al. 2012). At higher latitudes, the microwave emission vanishes, indicating low B (Dobler 2012a; Ade et al. 2012).

These magnetic fields pose two challenges to my model. First, the synchrotron losses cannot be much faster than IC, or else there is not enough wind power to explain the γ-ray emission. But for $B \lesssim 6$ μG, $f_{\rm IC} \gtrsim 0.5$.

Second, if the magnetic pressure is much greater than $P_{\rm ext}$, that could invalidate my calculations of R_s. For $B = 4\ \mu$G, the magnetic pressure is 5000 K cm^{-3}, with a similar pressure in CRs. But the wind ram pressure at 4 kpc from the CMZ is only 2000 K cm^{-3}. The tension is eased if the CMZ SFR was $\sim 0.2\ M_\odot$ yr^{-1} in the recent past.

3.3. *Ultra-high energy CRs from the shocks?*

The maximum CR energy is limited by the magnetic field in the shock, the size of the acceleration region, and the radiative losses the CRs experience. The sheer size of the Bubbles makes them excellent places for the acceleration of extremely high energy particles (c.f., Cheng et al. 2012).

Electrons are highly radiative particles, so their maximum energy is set by IC or synchrotron losses. They gain a fraction $\sim v_\infty/c$ of their energy per Larmor orbit, for an acceleration time of $t_{\rm acc} \approx 20E/(3v_\infty eB)$ that rises with energy (Gaisser 1990). By setting $t_{\rm acc}$ to the energy loss time from synchrotron and IC losses, I derive

$$E_e \lesssim \sqrt{\frac{9eBv_\infty}{80\sigma_T U c}} m_e c^2 \approx 0.6\,{\rm PeV} \left(\frac{B}{5\ \mu{\rm G}}\right)^{1/2} \left(\frac{U}{{\rm eV\ cm^{-3}}}\right)^{-1/2}. \tag{12}$$

U includes the sum of the CMB energy densities and the magnetic field energy densities; Klein-Nishina effects suppress losses off starlight for PeV electrons, but electrons still radiate IC emission by upscattering the CMB. These γ-rays cascade down to TeV energies due to $\gamma\gamma$ absorption within the Galaxy (Moskalenko et al. 2006). TeV and PeV electrons are a possible source of TeV γ-rays from the Fermi Bubbles.

For protons, the most important constraint on the maximum energy is that their Larmor radius cannot be larger than the acceleration region (Hillas 1984). Requiring that $E_p \lesssim R_{\rm max} ZeB$, I find

$$E_p \lesssim 5Z\ {\rm EeV} \left(\frac{R_{\rm max}}{1\ {\rm kpc}}\right) \left(\frac{B}{5\ \mu{\rm G}}\right). \tag{13}$$

It takes $t_{\rm acc} \approx 4$ Myr to reach these energies if $B = 5\ \mu$G.

Nuclei at these energies are subject to photopion and photodissociation losses off the far infrared radiation from the Milky Way. Photodissociation and photopionic γ-rays cascade down to $\sim$100 TeV as they propagate to Earth. Photopions can also decay into neutrinos, which could be detected at $\sim$10 PeV.

4. DISCUSSION

The starburst power source of this model is known, the wind's existence is fairly well established, and its termination shock is almost inevitable if the wind becomes supersonic. Because the injection sites are the termination shocks, there are no problems with cooling. IC emission does not require dense reservoirs of gas or long timescales, and is consistent with the wind's mechanical power. Finally, the shock could be a source of TeV IC emission, PeV neutrinos, and EeV CRs.

There are several problems that must be addressed. First, why does the e^- spectrum harden at low energies (Su et al. 2010; Ade et al. 2012)? This could be a cooling break, but only if the electrons are just a few Myr old. Yet the CMZ is likely older ($\gg$ 10 Myr), so older electrons are probably present on some level. Have the older electrons simply diffused away? Perhaps the CMZ fluctuates on timescales of $\sim$ 10 Myr, but is steady on longer timescales. Then, the Fermi Bubbles' IC emission represents the electrons from the most recent pulse of star formation. Either way, low energy IC and radio emission should be visible surrounding the Fermi Bubbles. Or is the break in the injection spectrum itself?

Second, why are the Bubbles elongated? This indicates the termination shock radius depends on angle, which could happen if the external pressure depends strongly on midplane distance instead of distance to the Galactic Center, or if power is concentrated along the axis of the outflow cone.

Third, this model cannot explain the possible γ-ray jets discovered by Su & Finkbeiner (2012) in the inner regions of the Bubbles, which could be from Sgr A* itself. The "jets" could instead be related to stellar clusters in the CMZ (c.f., Carretti et al. 2013); perhaps they are smaller cluster winds within the CMZ wind.

Finally, the flat γ-ray surface brightness of the Bubbles is inconsistent with IC emission from a thin shell around the shocks (Su et al. 2010). If there is turbulence within the wind bubble, second order acceleration (perhaps of the adiabatically cooled initial CRs) might be able to solve this problem, but the model proposed by Mertsch & Sarkar (2011) proposes that the shock is growing instead of steady-state; any turbulence must be generated before the shock itself. Alternatively, if the wind fluctuates on Myr timescales, internal shocks may develop and accelerate CRs within the Bubbles (c.f., Dorfi & Breitschwerdt 2012).

I thank Roland Crocker and Todd Thompson for their comments. I was supported by a Jansky Fellowship from the National Radio Astronomy Observatory. The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under cooperative agreement with the National Science Foundation.

REFERENCES

Ade, P. A. R., Aghanim, N., et al. 2012, arXiv:1208.5483
Aharonian, F., et al. 2006b, Nature, 439, 695
Beck, R., & Krause, M. 2005, Astronomische Nachrichten, 326, 414
Bell, A. R. 1978, MNRAS, 182, 443
Biermann, P. L., Becker, J. K., Caceres, G., et al. 2010, ApJ, 710, L53
Bland-Hawthorn, J., & Cohen, M. 2003, ApJ, 582, 246
Breitschwerdt, D., McKenzie, J. F., & Voelk, H. J. 1991, A&A, 245, 79
Carretti, E., Crocker, R. M., Staveley-Smith, L., et al. 2013, Nature, 493, 66
Cheng, K.-S., Chernyshov, D. O., Dogiel, V. A., Ko, C.-M., & Ip, W.-H. 2011, ApJ, 731, L17
Cheng, K.-S., Chernyshov, D. O., Dogiel, V. A., et al. 2012, ApJ, 746, 116
Chevalier, R. A., & Clegg, A. W. 1985, Nature, 317, 44 (CC85)
Crocker, R. M., Jones, D. I., Melia, F., Ott, J., & Protheroe, R. J. 2010, Nature, 463, 65
Crocker, R. M., Jones, D. I., Aharonian, F., Law, C. J., Melia, F., & Ott, J. 2011a, MNRAS, 411, L11
Crocker, R. M., Jones, D. I., Aharonian, F., Law, C. J., Melia, F., Oka, T., & Ott, J. 2011b, MNRAS, 413, 763
Crocker, R. M., & Aharonian, F. 2011, Physical Review Letters, 106, 101102
Crocker, R. M. 2012, MNRAS, 423, 3512
Dobler, G., Finkbeiner, D. P., Cholis, I., Slatyer, T., & Weiner, N. 2010, ApJ, 717, 825
Dobler, G. 2012a, ApJ, 750, 17
Dobler, G. 2012b, ApJ, 760, L8
Dorfi, E. A., & Breitschwerdt, D. 2012, A&A, 540, A77
Fang, T., Bullock, J., & Boylan-Kolchin, M. 2013, ApJ, 762, 20
Finkbeiner, D. P. 2004, ApJ, 614, 186
Gaisser, T. K. 1990, Cambridge and New York, Cambridge University Press, 1990, p. 158

Guo, F., & Mathews, W. G. 2012, ApJ, 756, 181

Heckman, T. M., Armus, L., & Miley, G. K. 1990, ApJS, 74, 833

Hillas, A. M. 1984, ARA&A, 22, 425

Hooper, D., & Slatyer, T. R. 2013, arXiv:1302.6589

Hsu, W.-H., Putman, M. E., Heitsch, F., et al. 2011, AJ, 141, 57

Jokipii, J. R., & Morfill, G. 1987, ApJ, 312, 170

Jones, D. I., Crocker, R. M., Reich, W., Ott, J., & Aharonian, F. A. 2012, ApJ, 747, L12

Launhardt, R., Zylka, R., & Mezger, P. G. 2002, A&A, 384, 112

Mayya, Y. D., Bressan, A., Rodríguez, M., Valdes, J. R., & Chavez, M. 2004, ApJ, 600, 188

McKee, C. F., & Ostriker, J. P. 1977, ApJ, 218, 148

Mertsch, P., & Sarkar, S. 2011, Physical Review Letters, 107, 091101

Molinari, S., Bally, J., Noriega-Crespo, A., et al. 2011, ApJ, 735, L33

Moskalenko, I. V., Porter, T. A., & Strong, A. W. 2006, ApJ, 640, L155

Murray, N., Quataert, E., & Thompson, T. A. 2010, ApJ, 709, 191

Savage, B. D., Sembach, K. R., Wakker, B. P., et al. 2003, ApJS, 146, 125

Sofue, Y. 2000, ApJ, 540, 224

Spitzer, L., Jr. 1956, ApJ, 124, 20

Strickland, D. K., & Heckman, T. M. 2009, ApJ, 697, 2030

Su, M., Slatyer, T. R., & Finkbeiner, D. P. 2010, ApJ, 724, 1044

Su, M., & Finkbeiner, D. P. 2012, ApJ, 753, 61

Sutherland, R. S., & Dopita, M. A. 1993, ApJS, 88, 253

Völk, H. J., Aharonian, F. A., & Breitschwerdt, D. 1996, Space Sci. Rev., 75, 279

Völk, H. J., & Zirakashvili, V. N. 2004, A&A, 417, 807

Wolfire, M. G., McKee, C. F., Hollenbach, D., & Tielens, A. G. G. M. 1995, ApJ, 453, 673

Yao, Y., & Wang, Q. D. 2005, ApJ, 624, 751

Yusef-Zadeh, F., Hewitt, J. W., Arendt, R. G., et al. 2009, ApJ, 702, 178

Zubovas, K., King, A. R., & Nayakshin, S. 2011, MNRAS, 415, L21